



Head-to-head comparison of novel suction technologies in endourology: suction mini-PCNL versus flexible ureteroscopy using flexible and navigable suction ureteral access sheaths for renal stone management. An EAU-endourology, propensity score-matched analysis of 1372 patients

Angelo Cormio^{1,2} · Daniele Castellani^{2,3} · Steffi Kar-Kei Yuen^{3,4} · Khi Yung Fong⁵ · Nariman Gadzhiev⁶ · Jaisukh Kalathia⁷ · Amish Mehta⁸ · Vigen Malkhasyan⁹ · Nitesh Kumar¹⁰ · Edgar Beltrán-Suárez¹¹ · Abhay Mahajan¹² · Karl Tan¹³ · Kremena Petkova¹⁴ · Kazumi Taguchi¹⁵ · Mohamed Amine Lakmichi¹⁶ · Yiloren Tanidir¹⁷ · Marcos Cepeda^{3,18} · Alexey Martov¹⁹ · Zelimkhan Tokhtiyev¹⁹ · Lazaros Tzelves²⁰ · Andreas Skolarikos^{3,20} · Mahmoud Laymon²¹ · Esteban Acuña²² · Wissam Kamal²³ · Heng Chin Tiong⁴ · Leonardo Gomes Lopes²⁴ · Dmitriy Gorelov²⁵ · Bhaskar K. Somani^{3,26} · Thomas R. W. Herrmann^{3,27,28,29} · Vineet Gauhar^{3,30,31}

Received: 25 July 2025 / Accepted: 17 October 2025

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2025

Abstract

Background To compare the two suction technologies, namely suction mini-percutaneous nephrolithotomy (SM-PCNL) and flexible ureteroscopy using flexible and navigable suction ureteral access sheaths (FANS) for renal stone management.

Methods Propensity-matched pair analysis was conducted on prospectively collected data from two global registries. The Suction mini-PCNL registry (STUMPS) included 30 centers across 21 countries, and the Flexible and Navigable Suction Access Sheath in Ureteroscopy registry (FANS) involved 25 centers across 20 countries. Each group included 686 patients after matching for age, sex, Guy's stone score, and stone volume. The primary outcome was the stone-free rate (SFR), assessed by 30-day non-contrast CT scan, categorized as Grade A (100% clearance or zero residual fragment, ZRF), Grade B (single ≤ 4 mm residual fragment), or Grade C (any fragment > 4 mm or multiple fragments).

Results In the matched cohort, immediate ZRF was significantly higher in SM-PCNL (85.7% vs. 60.1%, $p < 0.001$), and reinterventions were fewer (0.9% vs. 4.4%, $p < 0.001$). FANS demonstrated significantly superior recovery profiles with lower postoperative pain scores (VAS 1.0 vs. 2.0, $p < 0.001$), shorter hospital stays (1.0 vs. 2.0 days, $p < 0.001$), and reduced need for basket extraction (9.5% vs. 24.5%, $p < 0.001$). Operative time difference of 6 min, favoring SM-PCNL was noted (46 vs. 40 min, $p < 0.001$). Complication rates were low and comparable: fever managed by observation (2.1% FANS vs. 3.2% SM-PCNL, $p = 0.29$) or antibiotics (3.1% FANS vs. 6.0% SM-PCNL, $p = 0.08$), blood transfusion (0.1% FANS vs. 0.3% SM-PCNL, $p > 0.99$). No sepsis case was reported for both modalities. Multivariable analysis identified SM-PCNL as a significant independent predictor of Grade A stone-free status (OR 4.06, 95% CI 3.06–5.43, $p < 0.001$). Notably higher stone complexity GSS 2 and 3 had significantly better predictive odds of lower SFR than stone volume. Stone volume was a predictor of any complications (OR 1.16, 95% CI 1.03–1.30, $p = 0.01$).

Conclusions Both techniques are safe and effective; SM-PCNL is better for achieving 100% stone clearance, especially in GSS 2 and 3 cases. FANS offers faster recovery and lower postoperative pain scores. The Guy's stone score is useful to predict 100% SFR for both modalities. Even though the proposed re-intervention rate was higher in FANS, as ours is only a 30-day follow-up study, the true reintervention rate and outcomes are lacking.

Keywords Kidney stones · Percutaneous nephrolithotripsy · Flexible ureteroscopy · Suction · Vacuum

Introduction

Nephrolithiasis affects up to 10% of the global population, with rising incidences due to dietary and environmental factors [1]. Over the past three decades, kidney stone disease (KSD) treatment has evolved from open surgery to minimally invasive techniques like percutaneous nephrolithotomy (PCNL) and retrograde intrarenal surgery (RIRS). A tailored approach is preferred based on stone size, location, patient anatomy, and clinical case complexity, such as congenital kidney anomalies and obesity [2–4]. Standard PCNL remains preferred for large or complex stones [5], with mini-PCNL reducing morbidity whilst maintaining high clearance rates [6, 7]. Suction integration in PCNL helps improve visibility, fragment evacuation, reduce operative time, hospital stays, and complications [8]. Technological advancements in RIRS include miniaturized scopes and flexible suction ureteral access sheaths, enhancing irrigation-aspiration techniques that are capable of reducing infection risk and improving fragment extraction [9, 10]. The use of mini-PCNL (M-PCNL) and FANS in stones 2–3 cm has shown that the latter is non-inferior to Mini-PCNL [11]. The study compared outcomes using the largest stone diameter and M-PCNL with M-PCNL sheaths of 18fr only with FANS, paving the way for using FANS in larger stone burdens. However, to our knowledge, there is no head-to-head comparison of two modalities, namely suction M-PCNL (SM-PCNL) and RIRS using FANS, both incorporating suction-aspiration for any renal stone burden, objectively assessed using the Guy's stone score (GSS) [12] and stone volume (SV) as two parameters for stone burden and complexity. The primary aim of this study is to report which of these are better able to achieve a 100% stone-free rate (SFR) and 30-day morbidity due to the same.

Methods

The Suction Technology Utility in Mini-PCNL Study (STUMPS) registry included patients diagnosed exclusively with kidney stones who were prospectively recruited from 30 medical centers across 21 countries between March and November 2024. For this study, only adult patients aged 18 years or older, with only kidney stones, and those with data reporting stone burden using the GSS and SV were included. The original details of all patients in this database were already published [13]. Patients under 18 years of age, with concurrent ureteral stones and a lack of data on both GSS and SV were excluded. GSS was calculated by the treating urologist.

In the global multicenter FANS study, patients were prospectively enrolled between August 1, 2023, to August 30,

2024, involving 25 medical centers in 20 Countries, all of whom participated in the STUMPS registry. Eligible participants were adults aged 18 years or older who successfully underwent RIRS with FANS using 10/12 Fr or 11/13 Fr sheaths for any stone burden, recorded using SV calculated and GSS. Pediatric patients and those with ureteral stones or patients who had incomplete data records were excluded. Institutional board review approval was obtained by the leading center (Asian Institute of Nephro-Urology, AINU #01/2024) for STUMPS and FANS (AINU-EC/28/2023) study. Inclusion criteria adapted from the global FANS study published earlier [14] included adult patients with normal renal anatomy who were deemed suitable to undergo RIRS using FANS-UAS for single or multiple renal stones in any location within the collecting system. Exclusion criteria included pediatric patients, anatomical renal abnormalities, concomitant lithotripsy for ureteral stone, and patients who lacked. Stone burden was reported using GSS and SV in these patients from the pre op NCCT scans. Baseline and operative characteristics were collected from electronic medical records. All patients had a non-contrast CT scan performed 6weeks before surgery to assess stone burden and renal complexity, and SFR was evaluated at 30 days after the procedure. SV (for the largest stone in cases of multiplicity) was calculated using the ellipsoid formula ($\text{length} \times \text{width} \times \text{depth} \times \pi \times 0.167$) and GSS was reported for all patients as per the original classification Surgeons performed suction mini-(SM)PCNL or RIRS procedures according to their standard clinical practices The choice of energy source, FANS-specific characteristics (sheath size, pre-stenting, and exit strategy), and PCNL-specific characteristics (patient positioning, puncture method, tract dilation method, exit strategy) were at the respective surgeons' discretion based on their experience and available resources.

The following FANS were employed: Clearpetra (Welllead Medical, Panyu, Guangzhou, China), Negative Pressure Ureteral Access Sheath (Innovex Medical, Shanghai, China), Elephant second generation (Zhejiang YiGao Medical Technology, Hangzhou, China), Suction UAS (SEPLOU, Beijing, China), Disposable Ureteral Guide Sheath (ZSR Biomedical Technology, Dongguan, Guangdong China China). For SM-PCNL, Clearpetra (Welllead Medical, Panyu, Guangzhou, China) and Shah sheath were used.

All positive preoperative cultures were treated as per the hospital's antibiogram. Anticoagulants or antiplatelets were stopped 3 days prior for all patients and restarted postoperatively at the surgeon's discretion. All patients received on-table single-dose antibiotic prophylaxis as per local practices, irrespective of prior treatment. Negative urine culture was not mandatory. Pain was evaluated by the treating surgeon on the first postoperative day using a 10-point visual

analog scale (VAS), with 1 representing the lowest level of pain [15]. No standardized pain protocol was employed and this was left to each participant center according to each case and local practice. All patients were followed up for 30 days. Complications were graded according to the Clavien classification [16]. The definition of sepsis adhered to the guidelines outlined in the Third International Consensus Definitions (sepsis-3) by the presence of at least two clinical criteria that constitute the quick SOFA score. Readmissions within 30 days for re-intervention or reported complications were documented. Stone-free status, as assessed by a non-contrasted CT scan, was categorized into one of the following categories:

- - *Grade A*: 100% stone-free, indicating zero residual fragments (ZRF),
- - *Grade B*: single RF measuring up to 4 mm in maximum diameter.
- - *Grade C*: single RF >4 mm in maximum diameter, or multiple RF of any size.

Operative time was defined as the interval between the start of cystoscopy and the completion of the procedure (i.e., stent placement, ureteral catheter placement, or no drainage). Ureterscopy time was defined as the duration from the insertion of the ureteral access sheath (UAS) to its removal. Laser lithotripsy time refers to the actual time spent using the laser for stone fragmentation, as recorded in the operative report or device log. Sheath accessibility to the renal collecting system was assessed intraoperatively in both the FANS and SM-PCNL groups to ensure effective access for stone extraction of the target calix or in case of fragments migration.

Statistical analysis

Continuous variables are described using median and inter-quartile range, while categorical variables are described using absolute numbers and percentages. To visualize the similarities and differences between both study arms, patient demographics, peri-operative parameters, and 30-day outcomes were compared between the groups using the χ^2 test for categorical parameters and the Mann-Whitney U test for continuous variables. Propensity score matching (PSM) was used to reduce confounding in the statistical comparisons. Propensity scores were calculated using a logistic regression model, and one-to-one nearest-neighbor matching for age, sex, GSS, and SV. To ensure optimal matching of covariates, the caliper width was started at 0.2 [17, 18], with the target of absolute standardized mean difference (ASMD) for all covariates of < 0.1 [18] being achieved at this starting

caliper width. All statistical comparisons were then repeated for the PSM cohort, like the overall cohort.

Finally, two multivariable logistic regression analyses were performed in the matched population to evaluate factors associated with Grade A stone-free status and all complications. These predictors are described using odds ratios (OR), 95% CI, and p-values. All statistical tests were performed using R Statistical language, version 4.3.0 (R Foundation for Statistical Computing, Vienna, Austria), with $p < 0.05$ indicating statistical significance.

Results

Table 1 shows that the unmatched cohort comprised 1,963 patients: 704 patients who underwent FANS (Group 1) and 1,259 patients who underwent SM-PCNL (Group 2). In the unmatched population, patients in both groups had similar median ages (50 years) but differed significantly in gender distribution (55.0% male in Group 1 vs. 40.2% male in Group 2, ASMD 0.30). The groups also showed significant differences in GSS distribution, with FANS patients having higher proportions of GSS 3 stones (13.8% vs. 8.4%, ASMD 0.25) and SM-PCNL patients having more GSS 1 stones (61.2% vs. 57.0%). SV was significantly larger in Group 2 (median 1.6 cm³ vs. 1.4 cm³, ASMD 0.31), and stone location patterns differed substantially between groups (ASMD 0.37). There were no GSS 4 in Group 1.

After propensity score matching, 686 patients were included in each group and successfully matched with excellent covariate balance (ASMD < 0.1 for all matched variables; Table 1). The energy source for lithotripsy after matching is presented in Fig. 1. The matched cohort demonstrated similar baseline characteristics between groups, with median ages of 50 years in Group 1 and 49 years in Group 2, comparable gender distribution (54.4% vs. 55.4% male), and well-balanced GSC, except GSS4 which was missing in FANS and only 1 patient noted in SM-PCNL group, and SVs. In the matched population, significant differences were observed in procedural parameters and outcomes between the groups (Table 2). Total operation time was longer for FANS procedures (median 46 min [IQR 37–61] vs. 40 min [IQR 27–66], $p < 0.001$). The need for basket extraction was significantly lower in Group 1 (9.5% vs. 24.5%, $p < 0.001$), while sheath accessibility to all intrarenal collecting system was comparable between groups (89.1% vs. 87.2%, $p = 0.34$).

Postoperative pain scores were significantly lower in Group 1 (median 1.0 [IQR 1.0–2.0] vs. 2.0 [IQR 1.0–3.0], $p < 0.001$). Regarding complications, there was no case of sepsis requiring intensive care unit admission (Clavien 4). The complication rate was low in both groups with a similar

Table 1 Baseline characteristics of unmatched and matched cohort

	Unmatched			Matched		
	FANS (<i>n</i> = 704)	SM-PCNL (<i>n</i> = 1259)	ASMD	FANS (<i>n</i> = 686)	SM-PCNL (<i>n</i> = 686)	ASMD
Age, years, median [IQR]	50 [36, 61]	50 [39, 60]	0.02	50 [36, 61]	49 [38, 60]	<0.01
Male gender, <i>n</i> (%)	387 (55.0)	506 (40.2)	0.30	373 (54.4)	380 (55.4)	0.02
ASA score, <i>n</i> (%)			0.11			0.14
1	368 (52.3)	692 (55.0)		360 (52.5)	373 (54.4)	
2	274 (38.9)	459 (36.4)		270 (39.3)	247 (36.0)	
3	62 (8.8)	103 (8.2)		56 (8.2)	61 (8.9)	
4	0	5 (0.4)		0 (0.0)	5 (0.7)	
BMI, kg/m ² , median [IQR]	26.0 [24.8, 29.4]	26.0 [23.0, 29.0]	0.08	26.0 [24.8, 29.4]	26.0 [23.0, 29.0]	0.19
First time stone former, <i>n</i> (%)	502 (71.3)	974 (77.4)	0.14	495 (72.2)	518 (75.5)	0.08
Right side stone, <i>n</i> (%)	284 (40.3)	614 (48.8)	0.17	272 (39.7)	336 (49.0)	0.19
Guy's stone score, <i>n</i> (%)			0.25			0.08
1	401 (57.0)	770 (61.2)		395 (57.6)	402 (58.6)	
2	206 (29.2)	365 (29.0)		200 (29.2)	206 (30.1)	
3	97 (13.8)	106 (8.4)		91 (13.3)	77 (11.2)	
4	0	18 (1.4)		0 (0.0)	1 (0.1)	
Hounsfield units, median [IQR]	1100 [976, 1220]	1191 [936, 1366]	0.25	1100 [967, 1215]	1200 [942, 1354]	0.27
Stone volume, cm ³ , median [IQR]	1.4 [0.8, 2.1]	1.6 [0.9, 3.1]	0.31	1.4 [0.8, 2.1]	1.4 [0.8, 2.1]	0.03
Stone location, <i>n</i> (%)			0.37			0.38
Multiple	25 (3.6)	0 (0.0)		25 (3.6)	0 (0.0)	
Upper pole	167 (23.7)	197 (15.6)		163 (23.8)	110 (16.1)	
Middle pole/pelvis	309 (43.9)	574 (45.6)		297 (43.3)	300 (43.7)	
Lower pole	203 (28.8)	488 (38.8)		201 (29.3)	276 (40.2)	

Bold values indicates that the results are statistically significant

FANS Flexible and navigable suction ureteral sheath, SM-PCNL suction mini-percutaneous nephrolithotripsy, BMI body mass index, ASMD absolute standardized mean difference, IQR interquartile range

incidence of blood transfusion (Clavien 2: 0.1% in Group 1 vs. 0.3% in Group 2, $p > 0.99$), postoperative fever ($> 38^{\circ}\text{C}$) managed by observation without antibiotics (Clavien 1: 0.1% in Group 1 vs. 0.3% in Group 2, $p > 0.99$), postoperative fever ($> 38^{\circ}\text{C}$) managed with more than 48 h of antibiotics (Clavien 2: 2.1% in Group 1 vs. 3.2% in Group 2, $p = 0.29$),

Hospital stay duration was significantly shorter for FANS procedures (median 1.0 day [IQR 0.0–1.0] vs. 2.0 days [IQR 2.0–3.0], $p < 0.001$). Readmission rates within 72 h were comparable between groups (1.7% vs. 1.2%, $p = 0.50$), but.

30-day CT scan demonstrated a significant advantage for ZRF for Group 2. Grade A stone-free status was achieved in 85.7% of SM-PCNL patients compared to 60.1% of FANS patients. A significantly higher planned reintervention for RF(s) after 30 days in Group 1 (4.4% vs. 0.9%, $p < 0.001$) was noted.

The multivariate analysis demonstrates that SM-PCNL (OR 4.06 95% CI 3.06–5.43, $p < 0.001$) was the only factor associated with higher odds of being grade A stone-free (Table 3). Higher GSS were associated with lower odds of ZRF (GSS 2: OR 0.60, 95% CI 0.44–0.81, $p = 0.001$; GSS 3: OR 0.36, 95% CI 0.24–0.56, $p < 0.001$). For complications,

only SV emerged as a significant predictor of increased odds (OR 1.16, 95% CI 1.03–1.30, $p = 0.01$).

Discussion

Recent technological innovations in both PCNL and RIRS, such as suction and vacuum devices, have emerged as a potential game changer in the management of renal stones [14, 19]. The foundational research from the global STUMPS and FANS registries proved that these technologies are feasible, safe, and have promising outcomes across various patient groups [13, 14]. A recent RCT [11] comparing non-suction M-PCNL with FANS also demonstrated that for stone diameters of 2–3 cm, FANS is non-inferior. There were no SM-PCNL cases, nor did it use the GSS to represent stone complexity. The excellent RCT was only done in high-volume centres with highly experienced surgeons in a well-protocolised environment, and thus excellent non-inferiority outcomes were demonstrated.

To the best of our knowledge, our prospective, multi-center comparative study represents the first head-to-head analysis comparing SM-PCNL and RIRS using FANS with a primary aim to achieve 100% stone-free status, assessing




Fig. 1 Energy sources for lithotripsy in unmatched and matched cohort

GSS and SV to represent the stone burden in a real-world practice using any device for either FANS or SM-PCNL [20]. As SV is better adopted in endourological management of KSD, 100% SFS or ZRF assessed at 30 days with an NCCT is considered the best measure for a successful intervention and is proposed as a better benchmark in KSD intervention [21]. Our findings demonstrate that SM-PCNL was significantly more effective in achieving complete stone clearance, whereas FANS offered advantages in terms of postoperative recovery and morbidity. Notably, both techniques were associated with low complication rates, confirming their overall safety in appropriately selected patients.

The superior 100% stone-free rate observed with SM-PCNL (85.7% vs. 60.1%) aligns with existing literature supporting the efficacy of miniaturized PCNL approaches. Previous studies have shown that mini-PCNL maintains comparable SFRs to standard PCNL while reducing morbidity [22, 23]. Even in the RCT [11], it was observed that the FANS group had a non-inferior immediate SFR (84% vs. 85%; 95% CI) and a significantly longer operative time of 29 min. We found that in our study, in the PSM group, the difference was 6 min more for Group 1 (Table 2, $p < 0.001$).

Multivariable regression analysis confirmed SM-PCNL as an independent predictor of ZRF (OR 4.06). Further, GSS 2 and 3 were independently associated with lower odds of achieving a Grade A SFS. These findings reinforce the prognostic utility of the GSS system, previously validated as a complexity-based risk stratification tool [12], and reinforce the importance of stone complexity scoring in treatment selection and outcome prediction. The planned higher reintervention rate following FANS (4.4% vs. 0.9%) must be balanced against the reduced morbidity and faster recovery associated with this approach, and a lack of longer follow-up limits our ability to actually report the true re-intervention. It has been reported that whilst an NCCT scan is the best way to predict and report RF, often the timing of the same is debatable, and that a 30-day NCCT scan may unnecessarily lead to increased re-intervention [24]. As a consequence, the choice between SM-PCNL and FANS should be carefully individualized based on patient factors, stone characteristics, and treatment priorities, but GSS should be considered instead of just SV or diameter if these two interventions are to be compared. SM-PCNL may be preferred when complete stone clearance is paramount, particularly for complex stones or patients. FANS may be

Table 2 Procedural and postoperative characteristics in unmatched and matched cohort

	Unmatched			Matched		
	FANS (n=704)	SM-PCNL (n=1259)	p	FANS (n=686)	SM-PCNL (n=686)	p
Total operation time, minutes, median [IQR]	46 [37, 60]	41 [29, 65]	<0.001	46 [37, 61]	40 [27, 66]	<0.001
Basket required for stone extraction, n (%)	66 (9.4)	295 (23.4)	<0.001	65 (9.5)	168 (24.5)	<0.001
Sheath able to access all of the renal collecting system, n (%)	627 (89.1)	990 (87.5)	0.36	611 (89.1)	567 (87.2)	0.34
Intraoperative stone-free status			<0.001			<0.001
100% clear	395 (56.1)	1082 (85.9)		388 (56.6)	591 (86.2)	
Only dust remains	237 (33.7)	156 (12.4)		226 (32.9)	83 (12.1)	
Fragments remain	72 (10.2)	21 (1.7)		72 (10.5)	12 (1.7)	
Collecting system injury requiring stent, n (%)	5 (0.7)	17 (1.4)	0.28	5 (0.7)	10 (1.5)	0.30
Postoperative fever (>38° C) managed by observation without antibiotics (Clavien grade 1), n (%)	14 (2.1)	32 (2.5)	0.62	14 (2.1)	22 (3.2)	0.29
Postoperative fever (>38 °C) managed with antibiotics in the ward (Clavien 2), n (%)	9 (2.9)	73 (5.8)	0.06	9 (3.1)	41 (6.0)	0.08
Blood transfusion (Clavien 2), n (%)	1 (0.1)	15 (1.2)	0.03	1 (0.1)	2 (0.3)	>0.99
Bleeding requiring multiple bladder washouts/irrigations (Clavien grade 2), n (%)	0	4 (0.3)	0.33	0	4 (0.6)	0.13
Postoperative pain score, median [IQR]	1.0 [1.0, 2.0]	2.0 [1.0, 3.0]	<0.001	1.0 [1.0, 2.0]	2.0 [1.0, 3.0]	<0.001
Stone-free status on 30-day CT scan, n (%)	420 (59.6)	1066 (84.7)	<0.001	412 (60.1)	588 (85.7)	<0.001
Grade A: zero RF	275 (39.1)	160 (12.7)		265 (38.6)	83 (12.1)	
Grade B: single RF ≤4 mm	9 (1.3)	33 (2.6)		9 (1.3)	15 (2.2)	
Grade C: single RF >4 mm/multiple RF of any size						
Hospital stay, days, median [IQR]	0.0 [0.0, 1.0]	2.0 [2.0, 3.0]	<0.001	1.0 [0.0, 1.0]	2.0 [2.0, 3.0]	<0.001
Readmission for any reason within 72 h, n (%)	14 (2.0)	17 (1.4)	0.37	12 (1.7)	8 (1.2)	0.50
Reintervention for RF(s) after 30 days, n (%)	30 (4.3)	15 (1.2)	<0.001	30 (4.4)	6 (0.9)	<0.001

Bold values indicates that the results are statistically significant

RF residual fragment, FANS Flexible and navigable suction ureteral sheath, SM-PCNL suction mini-percutaneous nephrolithotripsy, IQR interquartile range

Table 3 Multivariable regression analysis of factors affecting grade A stone-free status and overall complications

	Grade A stone-free status		All complications	
	OR (95%CI)	p	OR (95%CI)	p
Suction mini-PCNL (reference FANS)	4.06 (3.06–5.43)	<0.001	0.71 (0.49–1.04)	0.08
Stone volume	0.94 (0.85–1.03)	0.18	1.16 (1.03–1.30)	0.01
Guys' stone score (reference 1)				
2	0.60 (0.44–0.81)	0.001	1.15 (0.74–1.76)	0.53
3	0.36 (0.24–0.56)	<0.001	1.64 (0.94–2.79)	0.07
Hounsfield units	1.01 (0.98–1.04)	0.31	1.15 (0.74–1.76)	0.64

Bold values indicates that the results are statistically significant

FANS Flexible and navigable suction ureteral sheath, PCNL percutaneous nephrolithotripsy

more appropriate for patients prioritizing rapid recovery, those with comorbidities increasing surgical risk, or when the retrograde approach offers better access to specific stone locations. Even though counselling for a second stage possibility, especially in GSS2 and 3 cases, must be done, caution may be exercised to consider the modality of re-treatment. Notably, the FANS study also reported that even in normal kidneys with a median SV of 1.2 cm³, grade A SFS was achieved in 57.8% cases, yet the actual reintervention was only seen in 11 patients. Again, a larger stone burden (SV 3000 mm³ and above) was significantly associated with reduced 100% SFR and contributed to re-intervention.

While FANS demonstrated lower SFR compared to SM-PCNL in our study, the 60.1% Grade A stone-free rate, even with complex stones (GSS 2 and GSS 3), represents a significant improvement over traditional RIRS approaches. Recent international multicenter randomized trials have demonstrated that suction ureteral access sheaths achieve superior stone-free rates compared to traditional sheaths in RIRS, supporting the value of this technology. Cacciatore et al. found that the absence of fragments larger than 3 mm 30 days after RIRS was 95% in the FANS group as

compared with only 67% in the standard non-suction sheath [25]. More recently, a meta-analysis reiterated that FANS provided superior stone clearance and reduced complications compared to non-suction sheaths [26].

Beyond efficacy, the procedural and recovery profiles varied significantly between techniques. The FANS group needed fewer basket extractions (9.5% vs. 24.5%), showing that FANS efficiently evacuates fragments without the need for active retrieval. This could potentially reduce the costs of accessories [27]. Conversely, SM-PCNL required more manual basket retrieval despite active suction, likely due to stone migration or the need to clear large fragments.

Operative time difference of 6 min favoring SM-PCNL (46 vs. 40 min) aligns with previous comparative studies showing that RIRS usually involves longer operative times, particularly in cases of multiple or lower pole stones, due to complex deflection angles and limited outflow capacity [28, 29]. Whilst it was only 6 min, we are unable to quantify what might have been the exact reasons for the same. Likely, it is multifactorial influenced by the surgeon's experience, moving the FANS sheaths to various parts of the kidney for inspection, especially into the lower pole or complex anatomy, and final exit strategy deployed. Yet this is significantly less than the 29-minute difference reported by Zeng et al. [11]. Interestingly, in the RCT [11], the mean Operative Time (OT) in the M-PCNL group was 59 min, whilst in our study, OT for SM-PCNL was 40 min. Perhaps SM-PCNL can facilitate stone evacuation much faster and may be a better choice than M-PCNL alone when considering bigger stone burdens, and reiterates previous studies [30].

Additionally, SM-PCNL has been shown to reduce OT and intrarenal pressure (IRP), which may facilitate clearer vision and faster stone clearance [8]. We acknowledge that our study lacks IRP data to make deductions on which suction modality has a better IRP safety profile. It has, however, been postulated and well proven in studies that novel non-invasive monitoring techniques for IRP are available and have better SFR and complication outcomes in SM-PCNL and FANS [31–33].

Interestingly, mild intraoperative bleeding was more frequently reported in the FANS group (5.7% vs. 2.2%). While this did not obscure vision or affect procedural safety, it may reflect differences in intrarenal pressure dynamics and fluid handling between the two techniques. Prior studies have shown that mini-PCNL typically maintains lower IRP values, reducing the risk of pyelovenous backflow and infection-related complications [34]. Conversely, fURS may generate transient IRP peaks exceeding 100–400 cm H₂O unless actively controlled [34]. FANS technology, which incorporates suction, may help modulate IRP depending on the sheath design and irrigation settings. Maintaining IRP

below 30 cm H₂O is generally considered a critical safety threshold [34].

Regarding postoperative recovery, our findings demonstrated that patients treated with FANS reported significantly lower postoperative pain (median VAS 1.0 vs. 2.0) and shorter hospital stays (1 vs. 2 days), which represent important quality-of-life advantages. These findings align with the minimally invasive nature of retrograde approaches and the reduced tissue trauma compared to percutaneous procedures. These are consistent with the literature supporting FANS as a minimally invasive profile and outpatient feasibility, as previously indicated [11, 14, 35, 36].

Finally, in the multivariate model, SV emerged as an independent predictor of complications (OR 1.16), in line with previous literature linking increased stone burden with greater perioperative risk [37]. Larger SVs have been linked to longer operative times, increased bleeding, higher intrarenal pressure, and greater risk of postoperative infection or need for transfusion [38–40]. In particular, stone size has been shown to correlate with higher Clavien–Dindo complication grades following PCNL [7]. These findings highlight the importance of preoperative risk stratification in patients with complex stones and may support using SM-PCNL over FANS for its superior evacuation efficiency, particularly in those patients having large and complex stones. Increasingly, with the wider adoption of FANS, it's likely that more studies will evaluate SM-PCNL with FANS, GSS, and SV as significant predictors of ZRF and complications, respectively (Table 3). These may be the better variables to consider, as seen in our MVA.

Our study is not devoid of limitations. The non-randomized design, despite propensity score matching, may introduce selection bias. Nevertheless, both registries are prospective, large, and multicenter international, depicting a real-world practice. While this geographic and institutional diversity enhances generalizability, it inevitably introduces heterogeneity in terms of surgeon expertise, procedural technique, instrumentation, and perioperative care pathways. Although PSM was used to adjust for key confounders, unmeasured variables, such as surgical experience, center volume, and local protocols, may have influenced the outcomes. Moreover, the observational and non-randomized nature of the study prevents us from establishing causal relationships. Notably, the lack of detailed intraoperative data, like irrigation pressures and laser settings, limits understanding of the mechanisms driving outcome differences between the two techniques. Importantly, patient-reported outcomes such as quality of life, return to normal activities, or satisfaction were not collected, and these factors are increasingly valued in comparative surgical studies. This is undoubtedly a further study limitation. Additionally, follow-up was limited to 30 days; therefore, the actual re-interventions performed

and their findings are unavailable. Even though functional renal outcomes and long-term anatomical outcomes in different GSS groups are missing, it's well proven that using FANS in normal kidneys is anatomically and physiologically safe in adults [41, 42].

The use of GSS as a matching covariate between PCNL and RIRS groups may be inappropriate, as GSS was originally developed and validated specifically for PCNL procedures and reflects complexity factors unique to PCNL that do not translate to equivalent complexity in RIRS. However, GSS is a reasonable surrogate marker to standardize comparisons across groups in the absence of a universally accepted scoring system for RIRS as demonstrated by recent comparative studies which showed that GSS predict surgical results in RIRS as well, with SFRs decreasing and complications increasing as GSS increased for both procedures [43, 44]. Nevertheless, the significant baseline imbalance in lower calyceal stone distribution between groups (29.3% vs. 40.2%) represents another potential bias, as lower pole stones are traditionally considered more challenging for RIRS due to unfavorable ureteroscope angles and gravity-dependent drainage. Conversely, a recent study of RIRS with FANS comparing patients with lower pole stones versus other locations showed that the SFR of lower pole stones is comparable to non-lower pole locations, suggesting that the imbalance in lower pole stones might not have significantly biased our findings [45].

Finally, as this was a comparative study where the STUMPS registry had used 4 mm as a cutoff, this was also applied to FANS patients for statistical analysis. Perhaps if we used 2 mm as a cutoff threshold for RF, the outcome reporting may have been different. As highlighted by Zeng et al. [11], future randomized trials with standardized protocols and long-term follow-up are needed to refine the comparative roles of these suction modalities to enable a patient-tailored approach, also considering cost-effectiveness.

Conclusion

Although both techniques are safe and effective, SM-PCNL is better for achieving 100% stone clearance or ZRF, especially in GSS 2 and 3 cases. FANS offers faster recovery and lower postoperative pain scores. The GSS is useful to predict 100% SFR for both modalities. Even though the proposed re-intervention rate was higher in FANS, as ours is only a 30-day follow-up study, the true re-intervention rate and outcomes are lacking. As zero sepsis and absence of major complications make both modalities safe, surgeons should individualize treatment based on counselling and their expertise.

Author contributions Conception and design: Vineet Gauhar, Angelo Cormio, Daniele Castellani. Acquisition of data: Steffi Kar-Kei Yuen, Nariman Gadzhiev, Jaisukh Kalathia, Amish Mehta, Vigen Malkhasyan, Nitesh Kumar, Edgar Beltrán-Suárez, Abhay Mahaja, Karl Tan, Kremena Petkova, Kazumi Taguchi, Mohamed Amine Lakmichi, Yiloren Tanidir, Marcos Cepeda, Alexey Martov, Zelimkhan Tokhtiyev, Lazaros Tzelves, Andreas Skolarikos, Mahmoud Laymon, Esteban Acuña, Wissam Kamal, Heng Chin Tiong, Leonardo Gomes Lopes, Dmitry Gorelov. Statistics: Khi Yung Fong. Drafting of the manuscript: Angelo Cormio, Vineet Gauhar, Daniele Castellani. Critical revision for important intellectual content: Bhaskar K. Somani, Thomas R. W. Herrmann. Supervision: Vineet Gauhar. All authors approved of the final version of the manuscript for submission.

Funding The authors did not receive support from any organization for the submitted work.

Data availability Data is available on request from the corresponding author.

Declarations

Conflict of interest The authors declare no conflict of interest.

Human rights Institutional board review approval was obtained by the leading center for STUMPS (Asian Institute of Nephro-Urology, AINU #01/2024) and FANS (AINU-EC/28/2023) study. The study was performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments.

References

1. Scales CD Jr., Smith AC, Hanley JM, Saigal CS (2012) Prevalence of kidney stones in the United States. *Eur Urol* 62:160–165
2. Türk C, Petřík A, Sarica K, Seitz C, Skolarikos A, Straub M et al (2016) EAU guidelines on interventional treatment for urolithiasis. *Eur Urol* 69:475–482
3. Cormio A, Mantovan M, Palantrani V, Beltrami M, Fuligni D, Passarella V et al (2025) A narrative review on extracorporeal shock wave lithotripsy, ureterolithotripsy, and percutaneous nephrolithotripsy in patients with anomalous kidneys. *Minerva Urol Nephrol* 77:43–51
4. Cormio A, Auciello M, Falagario UG, Ricapito A, Mangiatordi A, Castellani D et al (2024) Mini and standard percutaneous nephrolithotomy in obese patients. Results from a Single-surgeon large series. *Eur Urol Open Sci* 63:113–118
5. Quhal F, Seitz C (2021) Guideline of the guidelines: urolithiasis. *Curr Opin Urol* 31:125–129
6. Ruhayel Y, Tepeler A, Dabestani S, MacLennan S, Petřík A, Sarica K et al (2017) Tract sizes in miniaturized percutaneous nephrolithotomy: a systematic review from the European association of urology urolithiasis guidelines panel. *Eur Urol* 72:220–235
7. Sharma K, Sankhwar SN, Goel A, Singh V, Sharma P, Garg Y (2016) Factors predicting infectious complications following percutaneous nephrolithotomy. *Urol Ann* 8:434–438
8. De Stefano V, Castellani D, Somani BK, Giulioni C, Cormio A, Galosi AB et al (2024) Suction in percutaneous nephrolithotripsy: Evolution, Development, and outcomes from experimental and clinical studies. Results from a systematic review. *Eur Urol Focus*. 10: 154–68
9. Giulioni C, Castellani D, Traxer O, Gadzhiev N, Pirola GM, Tanidir Y et al (2024) Experimental and clinical applications and

- outcomes of using different forms of Suction in retrograde intrarenal surgery. Results from a systematic review. *Actas Urol Esp (Engl Ed)* 48:57–70
10. Pauchard F, Ventimiglia E, Corrales M, Traxer O (2022) A practical guide for intra-renal temperature and pressure management during rirs: what is the evidence telling Us. *J Clin Med* 11:3429
 11. Zeng G, Jiang K, Liu S, Wu R, Duan X, Chai CA et al (2025) Flexible ureteroscopy with a flexible and navigable suction ureteral access sheath versus mini-percutaneous nephrolithotomy for treatment of 2–3 cm renal stones: an international, multicenter, randomized, noninferiority trial. *Eur Urol*.
 12. Thomas K, Smith NC, Hegarty N, Glass JM (2011) The guy's stone score—grading the complexity of percutaneous nephrolithotomy procedures. *Urology* 78:277–281
 13. Gauhar V, Castellani D, Kalathia J, Mehta A, Gadzhiev N, Malkhasyan V et al (2025) Prospective multicenter real-world outcomes of Suction technology utility in Mini-PCNL study (STUMPS) in modern-day practice: formulation of the global STUMPS registry on behalf of the endourology section of the European association of urology and the Suction mini-PCNL collaborative study group. *World J Urol* 43:298
 14. Gauhar V, Traxer O, Castellani D, Sietz C, Chew BH, Fong KY et al (2024) Could use of a flexible and navigable Suction ureteral access sheath be a potential game-changer in retrograde intrarenal surgery? Outcomes at 30 days from a large, prospective, multicenter, real-world study by the European association of urology urolithiasis section. *Eur Urol Focus*. 10: 975–82
 15. Delgado DA, Lambert BS, Boutris N, McCulloch PC, Robbins AB, Moreno MR et al (2018) Validation of digital visual analog scale pain scoring with a traditional paper-based visual analog scale in adults. *J Am Acad Orthop Surg Glob Res Rev* 2:e088
 16. Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 240: 205–13
 17. Austin PC (2011) Optimal caliper widths for propensity-score matching when estimating differences in means and differences in proportions in observational studies. *Pharm Stat*. 10: 150–61
 18. Lunt M (2014) Selecting an appropriate caliper can be essential for achieving good balance with propensity score matching. *Am J Epidemiol* 179:226–235
 19. Jahrreiss V, Nedbal C, Castellani D, Gauhar V, Seitz C, Zeng G et al (2024) Is Suction the future of endourology? Overview from EAU section of urolithiasis. *Ther Adv Urol* 16:17562872241232275
 20. Zeng GH, Zhong W, Mazzon G, Zhu W, Lahme S, Khadgi S et al (2024) International alliance of urolithiasis (IAU) consensus on miniaturized percutaneous nephrolithotomy. *Mil Med Res* 11:70
 21. Tonyali S, Emiliani E, Şener TE, Pietropaolo A, Özsoy M, Aboumarzouk O et al (2022) Definition of clinically insignificant residual fragments after percutaneous nephrolithotomy among urologists: a world-wide survey by EAU-YAU endourology and urolithiasis working group. *Cent Eur J Urol* 75:311–316
 22. Ferakis N, Stavropoulos M (2015) Mini percutaneous nephrolithotomy in the treatment of renal and upper ureteral stones: lessons learned from a review of the literature. *Urol Ann* 7:141–148
 23. Liu Y, Zhu W, Zeng G (2021) Percutaneous nephrolithotomy with suction: is this the future? *Curr Opin Urol* 31:95–101
 24. Gauhar V, Castellani D, Chew BH, Smith D, Chai CA, Fong KY et al (2023) Does unenhanced computerized tomography as imaging standard post-retrograde intrarenal surgery paradoxically reduce stone-free rate and increase additional treatment for residual fragments? Outcomes from 5395 patients in the FLEXOR study by the TOWER group. *Ther Adv Urol* 15:17562872231198629
 25. Cacciatore L, Minore A, Bonanno L, Contessa P, Esperto F, Iannello AR et al (2025) Is flexible navigable suction ureteral access sheath (FANS) safer and more efficient than conventional sheaths? Italian multicentric experience. *World J Urol* 43:153
 26. Liu Q, Zeng T, Zhu S (2025) Flexible and navigable suction ureteral access sheath versus traditional ureteral access sheath for flexible ureteroscopy in renal and proximal ureteral stones: a meta-analysis of efficacy and safety. *BMC Urol* 25:127
 27. Lee YJ, Bak DJ, Chung J-W, Lee JN, Kim HT, Yoo ES et al (2016) Is it necessary to actively remove stone fragments during retrograde intrarenal surgery? *Investig Clin Urol* 57:274–279
 28. Geavlete P, Georgescu D, Niță G, Mirciulescu V, Cauni V (2006) Complications of 2735 retrograde semirigid ureteroscopy procedures: a single-center experience. *J Endourol* 20:179–185
 29. Lv G, Wang K, Zhang Z, Zhou C, Li Y, Zhang D (2022) Comparison of flexible ureteroscopy and mini-percutaneous nephrolithotomy in the treatment for renal calculi larger than 2 cm: a matched-pair analysis. *Urolithiasis* 50:501–507
 30. Li P, Ziyi H, Xia S, Tongxin Y, Guang W, Yongming J et al (2022) Comparison of vacuum suction sheath and non-vacuum suction sheath in minimally invasive percutaneous nephrolithotomy: a meta-analysis. *J Invest Surg* 35:1145–1152
 31. Bai J, Shangguan T, Zou G, Liu L, Xue X, Lin J et al (2024) Efficacy and intrarenal pressure analysis of flexible and navigable suction ureteral access sheaths with flexible ureteroscopy in modified surgical positions for 2–6 cm upper urinary tract stones: a multicenter retrospective study. *Front Med (Lausanne)* 11:1501464
 32. Yuen SKK, Somani B, Gauhar V (2025) Measuring, monitoring and reporting intrarenal pressure: a practical guide to endourologists from section of EAU endourology. *Curr Opin Urol* 35:399–411
 33. Zhu W, Yuen SKK, Cao J, Chai CA, Liu S, Du J et al (2025) Intrarenal pressure monitoring via flexible and navigable suction ureteral access sheath in retrograde intrarenal surgery: a preclinical animal study and a pilot clinical study. *Clin Transl Discov* 5:e70031
 34. Tokas T, Tzanaki E, Nagele U, Somani BK (2021) Role of intrarenal pressure in modern day endourology (Mini-PCNL and flexible URS): a systematic review of literature. *Curr Urol Rep* 22:52
 35. Koushik TPP, Meyyappan V, Aher NB, Sekar H, Thiruvengadam G, Krishnamoorthy S (2025) Miniperc percutaneous nephrolithotomy versus retrograde intrarenal surgery in the treatment of Juxta uretero-pelvic junction upper ureteric calculi: a prospective, randomized control study. *Urol Ann* 17:9–16
 36. Gauhar V, Somani BK, Heng CT, Gauhar V, Chew BH, Sarica K et al (2022) Technique, feasibility, utility, limitations, and future perspectives of a new technique of applying direct in-scope suction to improve outcomes of retrograde intrarenal surgery for stones. *J Clin Med* 11:5710
 37. Watanabe T, Komeya M, Odaka H, Kiuchi H, Saigusa Y, Makiyama K et al (2024) Ureteral stone volume and female gender predicts perioperative complications after complete ipsilateral upper urinary tract stone removal using flexible ureterorenoscopy. *Int Urol Nephrol* 56:1611–1616
 38. Akman T, Binbay M, Sari E, Yuruk E, Tepeler A, Akcay M et al (2011) Factors affecting bleeding during percutaneous nephrolithotomy: single surgeon experience. *J Endourol* 25:327–333
 39. de la Rosette JJ, Opondo D, Daels FP, Giusti G, Serrano A, Kandasami SV et al (2012) Categorisation of complications and validation of the clavien score for percutaneous nephrolithotomy. *Eur Urol*. 62: 246–55
 40. Labate G, Modi P, Timoney A, Cormio L, Zhang X, Louie M et al (2011) The percutaneous nephrolithotomy global study: classification of complications. *J Endourol* 25:1275–1280
 41. Candela L, Gauhar V, Somani B, Fong KY, Persaud S, Castellani D et al (2025) Acute kidney injury following retrograde intrarenal surgery (RIRS) with flexible and navigable Suction ureteral access sheath (FANS): results from a prospective multicenter study. *Minerva Urol Nephrol*. 77: 356–64

42. Fong KY, Yuen SKK, Somani BK, Malkhasyan V, Tanidir Y, Persaud S et al (2025) Assessment of outcomes and anatomical changes in the upper urinary tract following flexible ureteroscopy with a flexible and navigable Suction ureteral access sheath: 3-Month results from a multicenter study. *Urology* 199:35–41
43. He Q, Huang Q, Hou B, Hao Z (2024) Prediction of percutaneous nephrolithotomy outcomes and flexible ureteroscopy outcomes using nephrolithometry scoring systems. *Int Urol Nephrol* 56(5):1585–1593
44. Yıldızhan M, Balcı M, Asil E et al (2021) Comparison of percutaneous nephrolithotomy and retrograde intrarenal surgery outcomes for kidney stones larger than 2 cm from guy's stone scoring system perspective. *Int J Clin Pract* 75(12):e14956
45. Shrestha A, Traxer O, Seitz C et al (2024) Assessing flexible ureteroscopy outcomes for lower pole versus non lower pole stones

using the flexible and navigable suction ureteric access sheath: a prospective multicenter study by EAU endourology and PEARLS group. *World J Urol* 43(1):41

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Angelo Cormio^{1,2} · Daniele Castellani^{2,3} · Steffi Kar-Kei Yuen^{3,4} · Khi Yung Fong⁵ · Nariman Gadzhiev⁶ · Jaisukh Kalathia⁷ · Amish Mehta⁸ · Vigen Malkhasyan⁹ · Nitesh Kumar¹⁰ · Edgar Beltrán-Suárez¹¹ · Abhay Mahajan¹² · Karl Tan¹³ · Kremena Petkova¹⁴ · Kazumi Taguchi¹⁵ · Mohamed Amine Lakmichi¹⁶ · Yiloren Tanidir¹⁷ · Marcos Cepeda^{3,18} · Alexey Martov¹⁹ · Zelimkhan Tokhtiyev¹⁹ · Lazaros Tzelves²⁰ · Andreas Skolarikos^{3,20} · Mahmoud Laymon²¹ · Esteban Acuña²² · Wissam Kamal²³ · Heng Chin Tiong⁴ · Leonardo Gomes Lopes²⁴ · Dmitriy Gorelov²⁵ · Bhaskar K. Somani^{3,26} · Thomas R. W. Herrmann^{3,27,28,29} · Vineet Gauhar^{3,30,31}

✉ Daniele Castellani
castellanidaniele@gmail.com

¹ Department of Urology and Renal Transplantation, Policlinico Riuniti di Foggia, University of Foggia, Foggia, Italy

² Urology Unit, Azienda Ospedaliero-Universitaria delle Marche, Via Conca 71, 60126 Ancona, Italy

³ Endourology Section of the European Association of Urology, Arnhem, The Netherlands

⁴ Department of Surgery, S.H. Ho Urology Centre, The Chinese University of Hong Kong, Hong Kong, China

⁵ Department of Urology, Sengkang General Hospital, Singapore, Singapore

⁶ Department of Urology, Saint-Petersburg State University Hospital, Saint-Petersburg, Russia

⁷ Fortune Urology Clinic, Botad, Gujarat, India

⁸ B T Savani kidney hospital, Rajkot, Gujarat, India

⁹ Department of Urology, Moscow Urology Center, Botkin Hospital, Moscow, Russia

¹⁰ Department of Urology, Ford Hospital and Research Centre, Patna, Bihar, India

¹¹ Department of Urology, Specialty Hospital La Raza, National Medical Center of the Mexican Institute of Social Security, Mexico City, Mexico

¹² Department of urology, Sai Urology Hospital and MGM Medical College, Aurangabad, India

¹³ Department of Urology, Veterans Memorial Medical Center, Quezon City, Philippines

¹⁴ Department of Urology and Nephrology, Military Medical Academy, Sofia, Bulgaria

¹⁵ Department of Nephro-Urology, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan

¹⁶ Department of Urology, Faculty of Medicine, University Hospital Mohammed the VIth, Cadi Ayyad University, Marrakesh, Morocco

¹⁷ Department of Urology, Medicana Atasehir Hospital, Istanbul, Turkey

¹⁸ Department of Urology, Hospital Universitario Río Hortega, Valladolid, Spain

¹⁹ Department of Urology and Andrology, IPPE of A.I. Burnazyan SSC FMBC, FMBA of Russia, Moscow, Russia

²⁰ Second Department of Urology, Sismanogleio Hospital, National and Kapodistrian University of Athens, Athens, Greece

²¹ Department of Urology, Urology and Nephrology center, Mansoura University, Mansoura, Egypt

²² Department of Urology, Institution Hospital Provincial de Ovalle, Ovalle, Coquimbo, Chile

²³ Urology Unit, King Fahd General Hospital, Jeddah, Saudi Arabia

²⁴ Department of Urology, Hospital Orizonti and Santa Casa de Belo Horizonte, Belo Horizonte, Minas Gerais, Brazil

²⁵ Department of Urology, First Saint-Petersburg State Medical University, Saint-Petersburg, Russia

²⁶ Department of Urology, University Hospital Southampton NHS Trust, Southampton, UK

²⁷ Department of Urology, Spital Thurgau AG, Kantonsspital Frauenfeld, Frauenfeld, Switzerland

²⁸ Division of Urology, Department of Surgical Sciences, Stellenbosch University, Stellenbosch Western Cape, South Africa

²⁹ Hannover Medical School, Hannover, Germany

³⁰ Department of Urology, Ng Teng Fong General Hospital, Singapore, Singapore

³¹ Department of Urology, Asian Institute of Nephrology and Urology, Hyderabad, Telangana, India